

## Images in medicine

### A deceitful case of an inguinal adenophlegmon as a presenting complaint for underlying metastatic cutaneous squamous cell carcinoma



Atef Mejri<sup>1,&</sup>, Khaoula Arfaoui<sup>1</sup>

<sup>1</sup>General Surgery Department Jendouba Regional Hospital, Jendouba, Tunisia

<sup>&</sup>Auteur corresponding: Atef Mejri, General Surgery Department Jendouba Regional Hospital, Jendouba, Tunisia

Received: 19 Oct 2019 - Accepted: 01 Dec 2019 - Published: 13 Dec 2019

Domain: Dermatology, General surgery

Key words: Cutaneous squamous cell carcinoma, adenophlegmon, bowenoid papulosis

Images in medicine | Volume 1, Article 56, 13 Dec 2019 | 10.11604/pamj-cm.2019.1.56.20708

Available online at: <https://www.clinical-medicine.panafrican-med-journal.com/content/article/1/56/full>

© Atef Mejri et al PAMJ - Clinical Medicine (ISSN: 2707-2797). This is an Open Access article distributed under the terms of the Creative Commons Attribution International 4.0 License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited

#### Images in medicine

*Squamous cell carcinoma is the second most common primitive skin cancer known to be responsible for the majority of deaths from skin cancer excluding melanoma. Through this presentation, we illustrate an unusual case of squamous cell carcinoma revealed by a super infected lymph node metastasis mimicking an adenophlegmon. A 58-year-old patient presented to the emergency room with an inguinal bulge evolving for 3 months, gradually increasing in size, recently becoming painful. The clinical examination revealed a red warm and painful crural swelling. No inter-toes space intertrigo or any other lower-extremity infections were found. The blood tests showed a biological inflammatory syndrome: WBC=11000, CRP=119. An abdominal CT scan concluded to an inguinal adenophlegmon. We decided then to proceed to*

*surgery. In view of the suspicious tissular aspect found per operatively, a biopsy was performed. The histological examination led to the diagnosis of squamous cell carcinoma. This case strongly suggests that underlying malignancy must be considered in front of every adenophlegmon whithing patients having cutaneous HPV induced lesions (red and yellow circle) even if the clinical examination is negative for of an invasive tumor and the usual surgical procedure of incision and drainage must be prohibited. The primary tumor, as in the case of our patient, may develop on bowenoid papulosis (yellow circle), which is characterized by multiple, small skin colored to reddish-brown papules, primarily occurring on the genitalia of young adults. The right management in this case should be based on surgical lymph node dissection with or without adjuvant radiation therapy.*



**Figure 1:** a deceitful case of an inguinal adenophlegmon as a presenting complaint for underlying metastatic cutaneous squamous cell carcinoma: after surgery aspect